

SOME ECO-FRIENDLY BIOPESTICIDES FROM NAGPUR DISTRICT

MESHRAM REENA, ANASANE PRADNYA & CHATURVEDI ALKA

Post Graduate Teaching Department of Botany, Rashtrasant Tukadoji Maharaj Nagpur University, Nagpur, India

ABSTRACT

Our country is endowed with rich diversity of species of Asteraceae family; a number of species are known to have pesticide properties, which can be proved to be pest specific. The present paper is the survey of such plants having pesticide properties and can be used for further studies.

KEYWORDS: Biopesticide, Asteraceae Family & Nagpur

Received: Mar 27, 2017; **Accepted:** Apr 18, 2017; **Published:** May 06, 2017; **Paper Id.:** IJBRJUN20171

INTRODUCTION

Nagpur district is endowed with a variety of flora, which can be considered as the store house of plant protection chemicals like flavonoids, alkaloids, essential oil etc. Nowadays, there is a renewed interest in botanicals because of the adverse effects of chemical pesticides. Still, recently only the plants with toxic chemicals were considered as suitable for plant protection. The discovery of antifeedant, insect growth regulators and other behavior modifying chemicals in plants against insect pest, fit into the concept of insect pest management assume importance (F. Brisca, 2009).

However, synthetic insecticides have serious drawbacks such as pest resurgence and resistance, lethal effects on non-target organisms, the risk of users, contamination, food residues and environmental pollution (Tapondjou *et al.*, 2002). Biological control is the use of one or more types of beneficial organisms, usually called natural enemies, to reduce the numbers of another type of organism, the pest (Susan E. Rice Mahr *et al.*, 2001). In addition, the precautions necessary to work with traditional chemical insecticides (Fields *et al.*, 2001) and the poor storage facilities of a traditional farmers in developing countries, which are unsuitable for effective conventional chemical control (Tapondjou *et al.*, 2002) emphasize the necessity of new effective methods for insect pest control of crops.

As Asteraceae family is nested high in the angiosperm phylogeny Asterideae/Asterales. The family contains the largest number of described, accepted species, of any plant family, CA. 24,000, with estimates of the total number reaching 30,000. There are 1600-1700 genera distributed around the globe except for Antarctica. Assuming that there are 250,000-350,000 species of flowering plants, then out of every eight to twelve species is in Compositae (about 10%) (Vicki A. Funk *et al.* 2002). Asteraceae family having compounds like flavonoids, saponins, tannins, phenols, steroids, terpenoids, etc. In the present paper some common plants of Asteraceae family having biopesticide potential are discussed.

Ageratum conizoides L. (Sahadevi, Goat weed).

Habit: Herb.

Plant Distribution: Common throughout along roadsides, waste places and forest edges.

It is an annual, branching herb which grows up to 1m in height. The stem and leaves are covered with fine white hairs, the leaves are ovate and up to 7.5cm long. The flowers are purple to white, less than 6mm across and arranged in a terminal inflorescence. The fruits are achenes and easily dispersed. Because of its propagation, it becomes a weed and causes problems for farmers and ecologists. Seeds are positively photoblastic, and viability is often lost within 12 months (Tailor Chandra Shekhar and Goyal Anju, 2012).

Flowers and Fruits: July-February

Compounds and Biopesticide Properties:

Aerial parts of *Ageratum conizoides* contained 11 chromenes, nine methoxyflavones (like 5,6,8,3',4',5'-hexamethoxyflavone and 8-hydroxy-5,6,7,3',4',5'-hexamethoxyflavone), sitosterol, (+)-sesamin and caryophyllene oxide. Other components of this plant are alkaloids, flavonoids, benzofurans and terpenoids. The plant shows insecticidal properties.

Ageratum conizoides has bioactive activity that may have agricultural use as shown by several research investigations, in different countries. Pereira in 1929, cited by Jaccoud (1961), reported use of the leaves as an insect (moth) repellent (Tailor Chandra Shekhar, 2012). The insecticide activity may be the most important biological activity of this species (Harborne, 2010). Both the essential oil and as well as the major components of the oil, namely the precocenes, have been reported to have antijuvenile hormonal activity (Shirwaikar A *et al.*, 2003).

A. Coincides possesses hormone antagonist precocene I and II and *Artemesia vulgaris* contain a substance sesquiterpenoid location named as 'Vaughan' (Sharada Singh and Rao P.J., 1999).

Hexane crude extract of *A. conizoides* (Coumarin) shows insecticide activity (toxicity) to *Sitophilous zeamais*, *Rhyzopertha dominica*, *Oryzaephilus surinamensis* of stored products (Marcio Dionizio Moreira *et al.*, 2007).

The extracts of *A. conyzoides* shows the larvicidal and ovicidal activity of aqueous and ethanolic extracts of leaves on the eggs (unembryonated and embryonated), first and second larval stages of *Heligmosomoides bakeri* (J. Wabo Pan *et al.*, 2011).

Caesulia axillaris Roxb. (Bangra, Caesulia)

Habit: Herb

Plant Distribution: It is a common weed in rice fields (Marshy places) throughout India.

Herbs, prostrate or suberect, annual. Leaves sessile, oblong-lanceolate, acute at apex. Heads compound, 1cm across, globose, white or purplish. Pappus of 2 ovate scales. Achenes 0.3*0.2 cm, flat, obovate, slightly notched.

Flowers and Fruits: September-February

Biopesticide Properties

Its essential oil has been reported against some insect pests causing deterioration of food commodities (Verma and Dubey, 2001). It can be used as a fungicide, the efficacy of *C. Auxiliaries* essential oil against *A. Flavours* LHPtc and other molds causing deterioration of naturally stored herbal raw materials so that the oil may be recommended against the fungi associated with raw during their processing (Prashant Kumar Mishra *et al.*, 2012).

The essential oil of *Caesulia auxiliaries* protected stored gram and wheat from *C. chinensis*, *S. oryzae* and *T. castaneum* for first 12 months of storage (Kumar R. *et al.*, 2007), (Verma J.*et al.*, 2001).

Bidens biternata (Lour.)

Habit: Herb.

Plant Distribution: throughout as a Common weed in gardens and in waste places.

Annuals herbs, up to 80cm high, erect; stem quadrangular, ribbed, hairy. Leaves alternate or imparipinnate to bipinnate, ultimate leaflets ovate, acute. Heads yellow c 1.3cm in diameter, in terminal or leaf opposed peduncles. Pappus setae 2-4 retrorsely hispid, bristly. Achenes c 1.5 cm long, linear, black.

Flowers and Fruits: August-September

Compounds and Biopesticide Properties

Heads contain polyacetylenes, chalcone and aurone glycosides, volatile oil, sterols, acylated okanin glycoside, gallic acid and phenols. Seeds and the whole plant are insecticidal. They are used for aphids, cutworm and termites.

It is used to repel pests in stored grain as well as aphids, ants, beetle, cabbage root fly, caterpillars, crickets, mites and termites. *Bidens biternata* also have some active components like chalcones, aurone glycosides, volatile oils, acylated okanin glycoside, gallic and oxalic acid and phenolic astringents. Some of these, particularly the volatile oils, are considered responsible for the mosquito repellent (T. C. Kazembe and S. Nkomo, 2012).

Leaves of *Bidens pilosa* and soap powder were used as an insecticide for the control of leaf miners and other insects. Similarly, PAN (2008) indicated that *B. pilosa* was effective against aphid, cutworm and termites. Likely, in this study, *B. pilosa* was minimized the number of onion thrips population, but exhibited low mortality rate percentage (Tadele Shiberu and Amin Mahammed, 2014).

Blumea eriantha DC.

Habit: Herb

Plant Distribution: Common throughout in fields and in open forests.

Herbs, erect, perennial. Leaves 2.0-7.5*1-2cm, lower leaves obovate, upper leaves elliptic-oblong, margins serrate. Heads yellow. Pappus white, longer than achenes, scanty. Achenes minutely hairy.

Flowers and Fruits: December-May

Compounds and Biopesticide Properties

The leaves yield an essential oil rich in ketones such as carvotanacetone and l-tetrahydrocarvone. Seeds also yield a volatile oil with a camphor-like smell. The essential oil is found larvicidal against *Culex pipiens* and *Culex fatigans*. It is generally used as an insect repellent.

Blumea eriantha is used as a mosquito repellent.

Echinops echinatus Roxb.

Habit: Herb

Plant Distribution: Common throughout in dry places under open situations.

Herbs, 30-90 cm high, much branched, rigid, annual, 30-90cm high. Leaves sessile, 2-10cm long, oblong, deeply pinnatifid, glabrous or minutely pubescent. Heads c 1.5 cm across (excluding spines), white, surrounded by strong white bristles. Pappus short, yellow, forms a short cylindrical tuft above achenes. Achenes 0.5 cm long, obconic, densely villous.

Flowers and Fruits: September-January

Compounds and Biopesticide Properties

Plant leaf extract of *Echinops echinatus* on mycelial growth of *S. rolfsii* were significant (Muhammad Farooq *et al.*, 2010). It has a potential of fungicide.

Eclipta prostrata (L.)

Habit: Herb.

Plant Distribution: Common throughout in wet situations.

Annual herbs, erect or prostrate. Leaves sessile, oblong-lanceolate or oblong-elliptic, sparsely strigose on both surfaces. Heads c 0.5 cm across, solitary or 2-together, on unequal, axillary peduncles, yellow. Achenes oblong-obovate, tubercles all over, trigonous, brown to black.

Flowers and Fruits: July- February

Compounds and Biopesticide Properties

The extract of roots, shoot and the whole plant showed nematicidal activity against nematode species like *Meloidogyne graminicola*, *Meloidogyne incognita* and *Rolylenchulus* species due to the presence of thiophenes and thienyls.

Field testing of *E. Alba* for seed treatment of sorghum. *E. Alba* extract and Calthio C caused significant inhibition of *Leptosphaeria saccharin* Breda de Haan (syn. *Phoma sorghina* Sacc.) *in vitro*. In field trials, a stimulatory effect on seedling emergence and yield increases of 7 to 38% were observed in *E. Alba* and the pesticide treatments, as compared to no treatment (Elisabeth P. Zida, 2012).

It shows mosquito larvicidal activity against *Culex sitiens* species (Latha C. *Et al.*, 1999).

Leaf and root extract shows ovicidal to *S. cerealella* property against stored insects of rice (Prakash and Rao, 1997).

Tagetes erecta L.

Habit: Herb.

Plant Distribution: The 'African Marigold', a native of Mexico is grown abundantly in gardens (Cooke, op. cit.; Naik, op. cit.),

Herbs, 0.5-1.2 m high, stout, erect, annual. Leaves opposite or upper alternate, pinnately divided into lanceolate, serrate segments. Heads solitary, peduncled. Marginal florets with yellow, tubular corollas. Achenes linear-oblong, blackish. Pappus scales united.

Flowers and Fruits: September-October**Compounds and Biopesticide Properties**

The flower heads of this plant, yield a volatile oil consisting of targeting, limonene, valeric acid, acumen, etc. the Acetone extract as well as the steam distillate is found larvicidal, growth regulator and adulticidal against *Culex quinquefasciatus*, *Anopheles Stephenson* and *Aedes aegyti* this plant is used as interrupt with *Abelmoschus esculentus* Moench (*Hibiscus esculentus* Linn.) for controlling root infesting nematodes.

Almost all species of *Tagetes* (Flowers, leaves, roots) are found insecticidal, repellent, fungicidal and nematocidal. Aqueous extract is used for ants, aphids and grasshopper whereas the fermented extract is found better for rice blast and tomato blight.

The crude extracts and its fractions of the flower of *Tagetes erecta* Linn. Was tested for insecticidal activity against a stored product insect pest, *Tribolium castaneum* (Herbst). The chloroform fraction showed highest toxicity against both the larvae and adults of *Tribolium castaneum* followed by 50 petroleum ether fraction and ethanol extract (Farjana Nikkon, *et al.*, 2009).

Tagetes oil is a new biochemical pesticide active ingredient intended for use as an insecticide / acaricide for the control of mites, whiteflies, aphids, thrips, mealybugs, scales, and psylla on a variety of food crops (Biopesticides registration action document, 2012).

Xanthium indicum Koen.

Habit: Herb.

Plant Distribution: Common throughout as a weed along roadsides and in waste places.

Herbs, annual; stem rough with short hairs. Leaves broadly ovate, triangular, appressed hairy. Heads 0.4-0.8 cm across, green (young), in terminal and axillary spikes. Fruits c 2 cm long, having 2 erect mucronate beaks, covered with hooked prickles. Achenes c 1*0.4 cm, oblong or ovoid, compressed, black.

Flowers and Fruits: January-May**Compounds and Biopesticide Properties**

The aerial parts contain a volatile oil rich in sesquiterpene lactones, xanthinin and xanthatin and alkaloids. The seeds contain a fixed oil (35%). The leaves exhibit nematocidal activity against various *Meloidogyne* species.

Tridax procumbens L.

Habit: Herb.

Plant Distribution: Common throughout as a weed in waste places.

Annuals or perennials, erect or procumbent herbs, branched at the base. Leaves opposite, ovate or lanceolate, margins serrate to coarsely incised dentate or trilobed. Heads heterogeneous, produced on erect, retrorsely hirsute and sparsely glandular peduncles, 10-25 cm long; ray florets white, ligules mostly bilabiate, disc florets tubular-campanulate, yellow, 5-lobed. Pappus of many aristate bristles, unequal in length. Achenes c 0.3 cm long, narrowly obconical, blackish, terete or ribbed, sparsely scarious, truncate at apex.

Flowers and Fruits: throughout the Year

Compounds and Biopesticide Properties

Tridax procumbens contains alkaloids, carotenoids, flavonoids (catechins and flavones), fumaric acid, FL-sitosterol, saponins and tannins. It is richly endowed with carotenoids, saponins, oleanolic acid and ions like sodium, potassium and calcium. Luteolin, glucoluteolin, quercetin and isoquercetin have been reported from its flowers (Sneha mundada, 2010).

Tridax procumbens along with eighteen weeds were selected, and a pesticide was produced from them by extraction method. The produced biopesticide was tested in 5 agricultural fields against tea, paddy, maize and cabbage plants. The pesticide was highly effective against aphids, leaf folder disease, thrips, Japanese beetle, army worm and the cutworm caterpillar. The effect of the pesticide derived from the agricultural weeds was around 50 to 75% more than the conventional chemical based pesticide (Manish Krishna, 2013).

CONCLUSIONS

Knowledge of the toxic plants, their toxic principles and their biological activity is of paramount importance not only to enable them to be utilized as natural pest control agents and replace the commercial synthetic pesticides, but also to enable us to understand the nature of their toxicity to non-targeted animals. Over 2,000 plant species have been reported to possess pesticidal activity (Crosby, 1971; Chakraborty and Basu, 1997) out of about 2,500,000 angiosperms so far documented, only a fraction of which have been analyzed.

In Asteraceae, variations are enormous in which each subgroup specializes in one or more characteristics. Since evolution is never ending, this group represents different lines on which the evolution progressed and would progress.

This family is the most diverse family and its each and every part having noble biopesticidal activity. The work done till today on its biopesticide activities like insecticidal activity (toxicity), larvicidal activity, nematocidal activity, fungicidal activity, acaricide. Various species act against aphids, leaf folder disease, thrips, Japanese beetle, etc.

In future, there is huge room for research in the direction of more pesticide activities of the Asteraceae family and to elucidate the mechanism of action of the same.

REFERENCES

1. Harborne B J (2010). *Phytochemical Methods, A guide to Modern Techniques of Plant Analysis*. New Delhi (India): Springer.
2. *Biopesticides Registration Action Document* (2012). U.S. Environmental Protection Agency Office of Pesticide Programs Biopesticides and Pollution Prevention Division.
3. Chakraborty I and S Basu (1997). *Biopesticides: for our healthier tomorrow*. Everyman Science XXXII I: 16-20.
4. Crosby D G (1971). *Minor insecticides of plant origin*. In: *Naturally Occurring Insecticides*, edited by M Jacobson, DG Crosby (Dekker Marcel Inc., New York, USA) 171-39.
5. Daniel M (2009). *Taxonomy Evolution at work*. Vadodara, India: Narosa Publishing House Pvt. Ltd.
6. Elisabeth P Z, S Lund and B N James (2012). *Seed treatment with a binary pesticide and aqueous extract of Eclipta alba (L.) Hassk. for improving sorghum yield in Burkina Faso*. *Journal of Tropical Agriculture* 50: 1-7.

7. Brisca F R and K Sahayaraj (2009). Influence of botanicals in total head protein of *Spodoptera litura* (Fab.). *Journal of Biopesticides*, **2(1)**: 52-55.
8. N Farjana, M Rowshanul Habib, M Rezaul Karim, Zennat Fardousi, M. Motiur Rehman and M. Ekramul Haque (2009). Insecticidal activity of flower of *Tagetes castaneum* (Herbst). *Research Journal of Agriculture and Biological Sciences* **5(5)**: 748-753.
9. Fields P G, Y S Xie, and X Hou (2001). Repellent effect of pea (*Pisum sativum*) fractions against stored product insects. *Journal of Stored Products Research* **37**:359-370.
10. Funk V A (2002). The Importance of Herbaria. *Plant Science Bulletin* **49**: 94-95.
11. Iqbal MAFU, ShM Iqbal, R Afzal and A Rasool (2010). In-vitro evaluation of different plant extracts on mycelial growth of *Sclerotium rolfsii* the cause of root rot of sugar beet. *Mycopath* **8(2)**: 81-84.
12. Jaccoud R J S (1961). Contribuição para o estudo formacognóstico do *Ageratum conyzoides* L. *Rev. Bras. Farm.* **42(11/12)**: 177-97.
13. Krishna M, T Khemchandani and RR Balaji (2013). Extraction of a novel biopesticide obtained from agricultural weeds useful for medicinal plants. *Academic Journals* **7(30)**: 2236-2242.
14. Kumar R, M Srivastava and N K Dubey (2007). Evaluation of *Cymbopogon martinii* oil extract for control of post harvest insect deterioration in cereal and pulse. *J. Food Prot.* **70**: 172-178.
15. Latha C, P D Vijhayakumar, S Velayudhan and A Joseph (1999). Biological activity of indigenous plant extracts as mosquito larvicides. *Indian J Exptl Biol***37**: 206-8.
16. Mahr S E R, R A Cloyd, D L Mahr and C S Sadof (2001). *Biological control of insects and other pests of greenhouse crops*. 581 pp. North Central Regional Publication. Cooperative Extension of the University of Wisconsin, Madison.
17. Krishna M, T Khemchandani and R R Balaji (2013). Extraction of a novel biopesticide obtained from agricultural weeds useful for medicinal plants. *Academic Journals* **7**: 2236-2242.
18. Moreira M D, M C Picanço, L C Almeida Barbosa, R N C Guedes, M R Campos, G Silva and J C Martins (2007). Plant compounds insecticide activity against *Coleoptera* pests of stored products. *Pesq. agropec. bras., Brasília.* **42(7)**: 909-915.
19. Mishra P K, R Shukla, P Singh, B Prakash and N K Dubey (2012). Antifungal and antiaflatoxigenic efficacy of *Caesulia axillaris* Roxb. essential oil against fungi deteriorating some herbal raw materials, and its antioxidant activity. *Industrial Crops and Products*. **36**: 74-80.
20. Moreira M D , C Picanc, A Barbosa, C Guedes, M Campos, G A Silva and J C Martins (2007). Plant compound insecticide activity against *Coleoptera* pests of stored products. *Pesq. agropec.bras.* **42(7)**: 909-915.
21. Farooq M A, U Iqbal, S M Iqbal, R Afzal and A Rasool (2010). In-vitro evaluation of different plant extracts on mycelial growth of *sclerotium rolfsii* the cause of root rot of sugar beet *Mycopath* **8(2)**: 81-84.
22. Mundada S and Shivhare R (2010). Pharmacology of *Tridax procumbens* a Weed: Review. *International Journal of PharmTech Research* **2(2)**: 1391-1394.
23. Nikkon F, M R M Habib, M R M Karim, Z Ferdousi, M M Rahman and M E Haque (2009). Insecticidal Activity of Flower of *Tagetes erecta* L. against *Tribolium castaneum* (Herbst). *Research Journal of Agriculture and Biological Sciences* **5(5)**: 748-753.
24. Nkomo and T C Kazembe (2012). Use of *Blumea alata*, *Bidens pilosa* and *Chenopodium ambrosioides* as Mosquito Repellents

- and Mosquitocides. *Bulletin of Environment, Pharmacology and Life Sciences* **1**(7): 59 - 66.
25. PAN (2008). *How to grow crops without endosulfan: Field guide to non-chemical Pest Management in banana, cabbage and other crucifers, cassava, citrus, coffee, corn, cotton and other fiber crops, cowpea, eggplant, forage crops, forest trees, garlic, lettuce, mango, mungbean, onion, ornamentals, peanut, pepper, pigeon pea, oil crops, ornamentals, potato, rice, sesame, sorghum, soybean, squash and other cucurbits, string bean, sweet potato, tea, tomato, and wheat production.* Pesticide Action Net work (PAN), Hamburg, Germany.
 26. Pon'e W J, OF Tankoua, J Yondo and M C Komtangi (2011). *The in vitro Effects of Aqueous and Ethanolic Extracts of the Leaves of Ageratum conyzoides (Asteraceae) on Three Life Cycle Stages of the Parasitic Nematode Heligmosomoides bakeri (Nematoda: Heligmosomatidae).* *Veterinary Medicine International*: 5.
 27. Prakash A, J Rao and V Nandagopal (2008). *Future of Botanical Pesticides in rice, wheat, pulses and vegetables pest management.* *Journal of Biopesticides* **1**(2): 154 - 169.
 28. Prakash A. and Rao (1997). *Botanical pesticides in Agriculture.* Boca Raton, USA.: CRC Lewis Pubs.
 29. Mishra P K, R Shukla, P Singh, B Prakash, A Kedia and N K Dubey (2012). *Antifungal, anti-aflatoxigenic, and antioxidant efficacy of Jamrosa essential oil for preservation of herbal raw materials.* *International Biodeterioration & Biodegradation* **74**: 11-16.
 30. PROTA (2004) *Vegetables: Plant Resources of Tropical Africa*, Wageningen, Netherlands.
 31. Sahayaraj and F B Renuga F B (2009). *Influence of botanicals in total head protein of Spodoptera litura (Fab.).* *Journal of Biopesticides* **2**(1): 52-55.
 32. Singh S and P J Rao (1999). *Effect of Artemesia vulgaris on development and reproduction of Spodoptera litura and Schistocerca gregaria.* *Indian Journal of Entomology* **61** (4): 305 – 311.
 33. Shekhar TC and G Anju (2012). *A Comprehensive Review on Ageratum conyzoides Linn. (Goat weed)* *International Journal of Pharmaceutical and Phytopharmacological Research* **1**(6): 391-395.
 34. Shiberu T, M Negeri and T Selvaraj (2013). *Evaluation of some botanicals and Entomopathogenic Fungi for the control of Onion thrips (Thrips tabaci L.)* West Showa, Ethiopia. *Plant Pathology & Microbiology* **4**(1).
 35. Shirwaikar A, P M Bhilegoankar, S Malini and J S Kumar (2003). *The gastroprotective activity of the ethanol extract of Ageratum conyzoides.* *J.Ethnopharmacol.* **86**(1): 117-121.
 36. Singh S and P Rao (1999). *Effect of Artemesia vulgaris on development Spodoptera litura and Schistocera gregaria.* *Indian journal of Entomology* **61**(4): 305-311.
 37. Mundada S and R Shivhare (2010). *Pharmacology of Tridax procumbens a Weed: Review.* *International Journal of Pharm Tech Research* **2**(2): 1391-139.
 38. Shiberu T and A Mahammed (2014). *The Importance and Management Option of Onion thrips, Thrips tabaci (L.) (Thysanoptera: Thripidae) in Ethiopia: A Review.* *J Horticulture* **1**:2.
 39. Tailor C S and A Goyal (2012). *"A Comprehensive Review on Ageratum conyzoides Linn. Goat weed)"* *Int. journal of Pharmaceutical & Phytopharmacological research* **1**(6): 391-395.
 40. Tapondjou L A, C Adler, H Bouda and D A Fontem (2002). *S. Efficacy of powder and essential oil from Chenopodium ambrosioides leaves as post-harvest grain protectants against six-stored products beelless* *Journal of Stored Products Research* **38**: 395-402.

41. Ugemuge N. R. *Flora of Nagpur*. Nagpur, India, n.d.
42. Varma J and N K Dubey (2001). Efficacy of essential oils of *Caesulia axillaris* and *Mentha arvensis* against some storage pests causing biodeterioration of food commodities. *Int. J. Food Microbiol* **68**: 207-210.
43. Verma A, B Singh and K P Singh (2001). Vegetable production in NEH region. In: *National Seminar on Approaches for Increasing Agriculture Productivity in Hill & Mountain Eco System*, ICAR Research Complex for NEH Region (Oct. 18-20), Barapani, pp. 110-111.
44. Verma S and R S Dubey (2001). Effect of cadmium on soluble sugar and enzymes of their metabolism in rice. *Biol. Plant.* **44**: 117-123.
45. Wabo P J, J D Ngankam Ntemah, C F Bilong Bilong and M Mbida (2011). A comparative study of the ovicidal and larvicidal activities of aqueous and ethanolic extracts of pawpaw seeds *Carica papaya* (Caricaceae) on *Heligmosomoides bakeri*. *Asian Pacific Journal of Tropical Medicine* 447-450.

